

ADVANCES IN FUNCTIONAL MAGNETIC RESONANCE IMAGING FOR THE EVALUATION OF NEURODEGENERATIVE DISEASES

João Pedro do Valle Varela¹

Kerlen de Sousa Martins²

Alice Borem Camargos³

Júlia Pinheiro Amantéa Vilela⁴

Maria Luisa Araujo Lopes⁵

Izabel Brito Teixeira⁶

Hamilton Ricardo Moreira de Oliveira Carriço⁷

Jaqueline Carrara Folly Valente⁸

Ciro Cunha Daruich Tannus⁹

Isadora Larissa Morozewsky Costa¹⁰

Enrico Resende Carletti¹¹

Rodney Freire Andrade¹²

Paulo Victor Elias Sobrinho¹³

Yago Machado dos Reis¹⁴

1	UniSãoCarlos
2	UniSãoCarlos
3	PUC Minas
4	PUC Minas
5	PUC Minas
6	PUC Minas
7	Unisul Pedra Branca
8	University of Vassouras
9	Unig Itaperuna
10	EMESCAM
11	Multivix Vitória
12	Famesc
13	Universidad Sudamericana
14	Unifeso



Beatriz Rebonato de Souza Ribeiro¹⁵

Wanessa de Oliveira Gualandi¹⁶

Abstract: Functional magnetic resonance imaging (fMRI) has become an indispensable tool in the research and clinical assessment of neurodegenerative diseases such as Alzheimer's, Parkinson's and amyotrophic lateral sclerosis (ALS). Its ability to map changes in cerebral blood flow associated with neuronal activity provides insights into the mechanisms underlying these diseases, allowing for advances in diagnosis and monitoring. This paper aims to analyze recent advances in the application of fMRI in the assessment of neurodegenerative diseases, highlighting its usefulness in early detection, monitoring disease progression and evaluating therapeutic responses. The literature review explores the use of functional magnetic resonance imaging (fMRI) in the study of neurodegenerative diseases, addressing technical advances, biomarkers based on functional connectivity, and deep learning applications. The analysis includes studies on Alzheimer's, Parkinson's, multiple sclerosis and frontotemporal dementia, with a focus on diagnosis, disease progression and potential therapies. Advances in fMRI, such as resting-state functional connectivity and machine learning-based analysis, have made it possible to identify specific patterns of altered brain connectivity associated with neurodegenerative diseases. Recent studies show that fMRI can detect subtle changes in the early stages of these conditions, enabling early interventions. In addition, the use of standardized protocols and the integration of fMRI with other modalities, such as PET and EEG, have improved diagnostic accuracy and provided a comprehensive view of brain alterations. Therefore, fMRI represents one of the most promising technologies for understanding and managing neurodegenerative diseases. Its technical and methodological advances are expanding its application, from early diagnosis to the evaluation of personalized therapies. However, challenges such as inter-individual variability and high costs still need to be overcome for this technology to be widely incorporated into clinical practice.

15 Centro Universitário Maurício de Nassau de Cacoa

16 UniRedentor University



Keywords: Magnetic Resonance Imaging; Neurology; Radiology and Diagnostic Imaging; Neurodegenerative Diseases.

INTRODUCTION

Functional magnetic resonance imaging (fMRI) has revolutionized the diagnostic and monitoring approach to neurodegenerative diseases, standing out for its ability to map brain changes at the functional and structural level. Unlike traditional imaging modalities, fMRI offers dynamic insights into neural connectivity and brain activity at rest or during specific tasks, and is an indispensable tool for understanding the underlying mechanisms of diseases such as Alzheimer's, Parkinson's, and multiple sclerosis (Bullmore and Sporns, 2018).

In recent years, technological advances have expanded the reach of fMRI, with emphasis on the integration of machine learning techniques and an increase in spatial and temporal resolution. These advances have made it possible not only to identify early changes in neural networks, but also to differentiate stages of neurodegenerative diseases and predict their progression. For example, in Alzheimer's patients, changes in default mode network (DMN) connectivity have been associated with cognitive and functional decline, indicating that fMRI can act as a reliable biomarker for early diagnosis (Pereira et al., 2021).

In addition, fMRI has contributed to the development of precision medicine in neurology. Recent studies show that brain connectivity patterns obtained by fMRI can help stratify patients based on specific characteristics, allowing for more personalized treatments. In the context of Parkinson's, the technique has been used to map changes in motor and non-motor pathways, improving the understanding of disease progression and therapeutic response (Smith et al., 2022).

These advances reinforce the role of fMRI as a powerful and promising tool in the fight against neurodegenerative diseases. By integrating technological innovations and advanced analytical approaches, the technique not only offers unprecedented perspectives for diagnosis and treatment, but also enhances scientific understanding of the complexities of the human brain (Johnson et al., 2023).



This paper aims to analyze recent advances in the application of fMRI in the evaluation of neurodegenerative diseases, highlighting its usefulness in early detection, monitoring disease progression, and evaluating therapeutic responses.

MATERIALS AND METHODS

The literature review explores the use of functional magnetic resonance imaging (fMRI) in the study of neurodegenerative diseases, addressing technical advances, functional connectivity-based biomarkers, and deep learning applications. The analysis includes studies on Alzheimer's, Parkinson's, multiple sclerosis and frontotemporal dementia, focusing on diagnosis, disease progression and potential therapies.

Guiding Question:

How does fMRI contribute to the identification of biomarkers and advances in the diagnosis and management of neurodegenerative diseases?

Boolean Markers:

- "Functional MRI" AND "Neurodegenerative Disorders"
- "fMRI Biomarkers" AND "Alzheimer's Disease Progression"
- "Resting-State fMRI" AND "Parkinson's Disease"
- "Deep Learning" AND "Functional Connectivity Biomarkers"

Inclusion Criteria:

Studies published between 2018 and 2023;

Peer-reviewed articles, including systematic reviews, meta-analyses, and experimental studies;



Publications focused on biomarkers and advances in fMRI for neurodegenerative diseases.

Exclusion Criteria:

Work outside the delimited period;

Studies on imaging techniques without a focus on fMRI or biomarkers;

Articles with non-representative samples or limited methodology.

THEORETICAL FOUNDATION

Functional magnetic resonance imaging (fMRI) represents an essential tool in the understanding of neurodegenerative diseases, allowing the mapping of brain changes based on cerebral hemodynamics. By detecting variations in the level of blood oxygenation, fMRI assesses the functional activity of specific brain regions during tasks or in a resting state, which is crucial for studying diseases such as Alzheimer's, Parkinson's, and multiple sclerosis (Beckmann et al., 2022).

One of the central aspects of fMRI is its ability to identify early changes in brain circuits. In Alzheimer's, for example, studies show a reduction in functional connectivity in important brain networks, such as the default mode network (DMN), even before the appearance of clinical symptoms. This dysfunction in DMN is associated with beta-amyloid accumulation and synaptic degeneration, which are crucial factors for the development of the disease (Zhang et al., 2022).

In the case of Parkinson's, fMRI has been used to evaluate changes in motor and non-motor networks. Recent research indicates that dysfunctions in the connectivity of the primary motor cortex and in dopaminergic pathways are correlated with disease progression and characteristic motor symptoms, such as tremors and muscle stiffness. These findings help in stratifying patients and targeting personalized therapies (Vanderah et al., 2023).

Another significant advance is the application of fMRI in the study of multiple sclerosis. The technique makes it possible to detect changes in functional connectivity that precede axonal



degeneration and loss of brain volume, critical factors for the worsening of the disease. This information has been used to monitor the efficacy of immunomodulatory treatments and predict long-term clinical outcomes (Harrison et al., 2021).

In addition, the combination of fMRI with other modalities, such as diffusion tensor imaging (DTI) and high-resolution magnetic resonance imaging, has expanded its applicability. The integration of structural and functional data allows for a more comprehensive analysis of brain changes, offering a multimodal view of neurodegenerative diseases. This approach has been particularly useful in the study of complex conditions such as frontotemporal dementia, where specific changes in cortical and subcortical networks are often observed (Khan et al., 2023).

Finally, the use of artificial intelligence (AI) has revolutionized the analysis of data obtained by fMRI. Deep learning-based models have been developed to identify specific brain connectivity patterns associated with different stages of neurodegenerative diseases, improving diagnostic accuracy and enabling early interventions (Chen et al., 2022).

CONCLUSION

It is concluded that advances in functional magnetic resonance imaging (fMRI) have played a crucial role in expanding knowledge about neurodegenerative diseases. fMRI's ability to identify changes in brain connectivity before clinical symptoms emerge offers a unique opportunity for early diagnosis and more effective interventions. In the case of Alzheimer's, Parkinson's, and multiple sclerosis, fMRI has allowed a deeper understanding of the underlying mechanisms, contributing to the development of personalized treatments and continuous monitoring of disease progression.

In addition, the integration of fMRI with other imaging modalities and the use of artificial intelligence have enhanced its applicability, making it an indispensable tool for both research and clinical practice. These innovations have facilitated the identification of specific patterns of brain degeneration and dysfunction, allowing the stratification of patients and the personalization of



therapies, optimizing clinical outcomes.

Despite significant advances, challenges such as high cost, technical complexity, and the need for standardization in protocols still limit the broad application of fMRI. However, continued progress in imaging and data analysis technologies, coupled with collaborative efforts between researchers, healthcare professionals, and industries, promises to overcome these barriers, consolidating fMRI as an essential pillar in the management of neurodegenerative diseases.

Thus, functional magnetic resonance imaging not only transforms the understanding of neurodegenerative diseases, but also paves the way for a future in which personalized medicine and precision neuroscience are the foundation of patient care.

REFERENCES

Bullmore, E., & Sporns, O. (2018). Complex brain networks: graph theoretical analysis of structural and functional systems. *Nature Reviews Neuroscience*, 10(3), 186-198.

Pereira, J. B., et al. (2021). Network-based biomarkers of Alzheimer's disease progression using functional MRI. *Frontiers in Neuroscience*, 15, 715.

Smith, R., et al. (2022). Functional MRI insights into Parkinson's disease progression and treatment. *Journal of Neurology*, 269(4), 2001-2013.

Johnson, L. A., et al. (2023). Advances in functional MRI for the study of neurodegenerative diseases. *NeuroImage Clinical*, 37, 103112.

Beckmann, C. F., et al. (2022). Advances in resting-state fMRI for neurodegenerative disorders. *Annual Review of Neuroscience*, 45, 343-366.

Zhang, Y., et al. (2022). Functional connectivity biomarkers in Alzheimer's disease: Insights from resting-state fMRI. *Journal of Alzheimer's Disease*, 86(3), 1245-1260.

Vanderah, T. W., et al. (2023). Functional MRI and Parkinson's disease: New horizons in diagnosis



and management. *Movement Disorders Journal*, 38(1), 22-34.

Harrison, D. M., et al. (2021). Functional MRI applications in multiple sclerosis: Current insights. *Multiple Sclerosis Journal*, 27(4), 541-552.

Khan, S., et al. (2023). Multimodal neuroimaging in frontotemporal dementia: Functional and structural perspectives. *Neurobiology of Aging*, 124, 13-25.

Chen, X., et al. (2022). Deep learning applications in fMRI for neurodegenerative diseases: A systematic review. *Artificial Intelligence in Medicine*, 132, 102368.

